



Data User Guide

NRT AMSR2 Daily L3 25 km Tb and Sea Ice Concentration Polar Grids

Introduction

The GCOM-W1 near real-time (NRT) AMSR2 Daily Level 3 gridded 25 km sea ice product includes brightness temperatures (Tb) at 6.9 through 89.0 GHz and sea ice concentrations. Data are mapped to a polar stereographic grid at 25 km spatial resolution. Sea ice concentration and brightness temperatures include daily ascending averages, daily descending averages, and daily averages. NRT products are generated within 3 hours of the last observations in the file by the Land Atmosphere Near real-time Capability for EOS (LANCE) at the AMSR Science Investigator-led Processing System (AMSR SIPS), which is collocated with the Global Hydrology Resource Center (GHRC) Distributed Active Archive Center (DAAC).

Notice:

All LANCE AMSR2 data should be used with the understanding that these are preliminary products. Cross calibration with AMSR-E products has not been performed. As updates are made to the L1R data set, those changes will be reflected in this higher level product.

Citation

Markus, T., W. Meier, and J. C. Comiso. 2015. NRT AMSR2 Daily L3 25 km Tb and Sea Ice Concentration Polar Grids [indicate subset used]. Dataset available online, [<https://lance.nsstc.nasa.gov/amr2-science/data/level3/seaice25>] from NASA LANCE AMSR2 at the GHRC DAAC Huntsville, Alabama, U.S.A. doi: http://dx.doi.org/10.5067/AMSR2/A2_S125_NRT

Keywords:

Brightness temperature, sea ice, sea ice concentration

LANCE

The Land Atmosphere Near real-time Capability for EOS (LANCE) makes EOS data from MODIS, AIRS, MLS, OMI, AMSR2, and MISR available within three hours of satellite overpass to meet the timely needs of applications such as numerical weather and climate prediction; forecasting and monitoring natural hazards, ecological/invasive species, agriculture, and air quality; providing help with disaster relief; and homeland security. Please note that LANCE has a rolling archive life of ten days on the HTTPS server. Once ten days pass following the data acquisition date, users must use the standard products.

If data latency is not a primary concern, please consider using science quality standard products. Science products are created using the best available ancillary, calibration and ephemeris information. Science quality products are an internally consistent, well-calibrated record of the Earth's geophysical properties to support science. The AMSR2 standard science quality data products will be available from the NSIDC DAAC.

Instrument Description

The Advanced Microwave Scanning Radiometer 2 (AMSR2) instrument aboard the Global Change Observation Mission - Water 1 (GCOM-W1) provides global passive microwave measurements of terrestrial, oceanic, and atmospheric parameters for the investigation of global water and energy cycles. Both AMSR2 and GCOM-W1 are built and operated by Japan Exploration Agency (JAXA). Data from this instrument are ingested from JAXA into NASA's LANCE element at the AMSR SIPS to be processed with US AMSR Science Team members' algorithms.

The AMSR instruments improved upon the heritage of the Scanning Multichannel Microwave Radiometer (SMMR), Special Sensor Microwave/Imager (SSM/I) and Tropical Rainfall Measuring Mission (TRMM) Microwave Instrument (TMI) instruments. Major improvements over those instruments included channels spanning the 6.9 GHz to 89 GHz frequency range, and higher spatial resolution from the 1.6 m reflector. More information about AMSR2 can be found at http://global.jaxa.jp/projects/sat/gcom_w/.

Investigators

Thorsten Markus
Cryospheric Science Laboratory
NASA Goddard Space Flight Center
Thorsten.Markus@nasa.gov

Walter Meier
Cryospheric Science Laboratory
NASA Goddard Space Flight Center
walter.n.meier@nasa.gov

Josefino C. Comiso
 Cryospheric Science Laboratory
 NASA Goddard Space Flight Center
 josefino.c.comiso@nasa.gov

File Naming Convention

The data is formatted using the following file naming convention.

Data: AMSR_2_L3_Sealce25km_X##_yyyymmdd.he5

Browse: AMSR_2_L3_Sealce25km_X##_yyyymmdd_f_CON.png

QA Summary Files: AMSR_2_L3_Sealce25km_X##_yyyymmdd.qa

Table 1: File naming convention variables

Variable	Description
X	Product Maturity code (Refer to table 2)
##	Two-digit file version number
yyyy	Four-digit year
mm	Two-digit month
dd	Two-digit day
f	N = Northern Hemisphere, S = Southern Hemisphere
.he5	HDF-EOS5 format
.xml	Metadata file
.met	Metadata file
.png	Portable Network Graphics format
.qa	GPS Quality Assessment Data

As NRT data are received from JAXA, partial daily products are generated and identified with a product maturity code of "P" in the filename. Once all Level-1R inputs are available, the complete daily product contains product maturity code "R" (near real-time) in the filename. Incremental processing makes data available to the user as it is received, rather than at the end of the day. Table 2 outlines the product maturity code variables used in the file naming convention.

Table 2: Variable Values for Product Maturity Code

Variable	Description
P	Partial daily product
R	Near real-time

Data Format Description

Data are stored in HDF-EOS5 format and are available via HTTP from the EOSDIS LANCE system at <https://lance.nsstc.nasa.gov/amr2-science/data/level3/seaice25/> or <https://lance.itsc.uah.edu/amr2-science/data/level3/seaice25/>. Please refer to Table 3 for information on the dataset characteristics.

Table 3: Dataset Characteristics

Characteristic	Description
Platform	Global Change Observation Mission - Water 1 (GCOM-W1)
Instrument	Advanced Microwave Scanning Radiometer 2 (AMSR2)
Projection	Polar Stereographic Projection*
Spatial Coverage	North Polar Grid N: 90, S: 30.98, E: 180, W: -180 South Polar Grid N: -39.23, S: -90, E: 180, W: -180
Spatial Resolution	25 km x 25km
Temporal Coverage	Start date: 09-06-2015 Stop date: Ongoing
Temporal Resolution	Daily averages, daily ascending average, daily descending average
Parameter	Brightness temperature Sea ice concentration Sea Ice Concentration differences between Bootstrap Basic Algorithm (BBA) and Enhanced NASA Team (NT2)
Processing Level	Level 3
Data Format	HDF-EOS 5

*For more information on the polar stereographic projection used for this dataset, please refer to the Spatial Coverage section within the corresponding AMSR-E [NSIDC documentation](#).

Data Parameters

Each data file contains core metadata, product-specific attributes, and data fields in 16-byte integer format (Int16). Please note that a scale factor has been applied to the brightness temperature data. To obtain the brightness temperature in kelvins (K), multiple data values by 0.1. The valid range of brightness temperature is approximately 50 to 300 K. Please refer to Table 4 and Table 5 for additional parameter information.

Table 4: Northern Polar Grids

Field Name	Description	Data Type	Unit	Scale Factor
SI_25km_NH_06H_ASC	6.9 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_06H_DAY	6.9 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_06H_DSC	6.9 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_06V_ASC	6.9 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_06V_DAY	6.9 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_06V_DSC	6.9 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_10H_ASC	10.7 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1

SI_25km_NH_10H_DAY	10.7 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_10H_DSC	10.7 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_10V_ASC	10.7 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_10V_DAY	10.7 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_10V_DSC	10.7 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_18H_ASC	18.7 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_18H_DAY	18.7 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_18H_DSC	18.7 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_18V_ASC	18.7 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_18V_DAY	18.7 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_18V_DSC	18.7 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_23H_ASC	23.8 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_23H_DAY	23.8 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_23H_DSC	23.8 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_23V_ASC	23.8 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_23V_DAY	23.8 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_23V_DSC	23.8 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_36H_ASC	36.5 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_36H_DAY	36.5 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_36H_DSC	36.5 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_36V_ASC	36.5 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_36V_DAY	36.5 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_36V_DSC	36.5 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_89H_ASC	89.0 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_89H_DAY	89.0 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_89H_DSC	89.0 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_NH_89V_ASC	89.0 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1

SI_25km_NH_89V_DAY	89.0 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1	
SI_25km_NH_89V_DSC	89.0 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_NH_ICECON_ASC	Sea ice concentration daily ascending average	Int16	N/A	N/A	
	Value				Description
	0				Open Water
	1 to 100				Percent Ice Concentration
120	Land Mask				
SI_25km_NH_ICECON_DAY	Sea ice concentration daily average	Int16	N/A	N/A	
	Value				Description
	0				Open Water
	1 to 100				Percent Ice Concentration
120	Land Mask				
SI_25km_NH_ICECON_DSC	Sea ice concentration daily descending average	Int16	N/A	N/A	
	Value				Description
	0				Open Water
	1 to 100				Percent Ice Concentration
120	Land Mask				
SI_25km_NH_ICEDIFF_ASC	Sea ice concentration daily ascending average using the difference between BBA and NT2 algorithms	Int16	N/A	N/A	
	Value				Description
	0				Open Water
	1 to 100 or -1 to -100				Percent difference between algorithms*
	120				Land Mask
	200 to 300				Missing NT2 value (200+BBA)**
	-200 to -300				Missing BBA value (-200-NT2)**
-310	Out-of-range value in V12 data***				
SI_25km_NH_ICEDIFF_DAY	Sea ice concentration daily average using the difference between BBA and NT2 algorithms	Int16	N/A	N/A	
	Value				Description
	0				Open Water
	1 to 100 or -1 to -100				Percent difference between algorithms*
	120				Land Mask
	200 to 300				Missing NT2 value (200+BBA)**
	-200 to -300				Missing BBA value (-200-NT2)**
-310	Out-of-range value in V12 data***				

SI_25km_NH_ICEDIFF_DSC	Sea ice concentration daily descending average using the difference between BBA and NT2 algorithms		Int16	N/A	N/A
	Value	Description			
	0	Open Water			
	1 to 100 or -1 to -100	Percent difference between algorithms*			
	120	Land Mask			
	200 to 300	Missing NT2 value (200+BBA)**			
	-200 to -300	Missing BBA value (-200-NT2)**			
-310	Out-of-range value in V12 data***				

Table 5: Southern Polar Grids

Field Name	Description	Data Type	Unit	Scale Factor
SI_25km_SH_06H_ASC	6.9 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_06H_DAY	6.9 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_06H_DSC	6.9 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_06V_ASC	6.9 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_06V_DAY	6.9 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_06V_DSC	6.9 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_10H_ASC	10.7 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_10H_DAY	10.7 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_10H_DSC	10.7 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_10V_ASC	10.7 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_10V_DAY	10.7 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_10V_DSC	10.7 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_18H_ASC	18.7 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_18H_DAY	18.7 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_18H_DSC	18.7 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_18V_ASC	18.7 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1
SI_25km_SH_18V_DAY	18.7 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1

SI_25km_SH_18V_DSC	18.7 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_23H_ASC	23.8 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_23H_DAY	23.8 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_23H_DSC	23.8 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_23V_ASC	23.8 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_23V_DAY	23.8 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_23V_DSC	23.8 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_36H_ASC	36.5 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_36H_DAY	36.5 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_36H_DSC	36.5 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_36V_ASC	36.5 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_36V_DAY	36.5 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_36V_DSC	36.5 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_89H_ASC	89.0 GHz horizontal daily average ascending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_89H_DAY	89.0 GHz horizontal daily average Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_89H_DSC	89.0 GHz horizontal daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_89V_ASC	89.0 GHz vertical daily average ascending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_89V_DAY	89.0 GHz vertical daily average Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_89V_DSC	89.0 GHz vertical daily average descending Tbs	Int16	Kelvin (K)	0.1	
SI_25km_SH_ICECON_ASC	Sea ice concentration daily ascending average		Int16	N/A	N/A
	Value	Description			
	0	Open Water			
	1 to 100	Percent Ice Concentration			
	120	Land Mask			
SI_25km_SH_ICECON_DAY	Sea ice concentration daily average		Int16	N/A	N/A
	Value	Description			
	0	Open Water			
	1 to 100	Percent Ice Concentration			
	120	Land Mask			

SI_25km_SH_ICECON_DSC	Sea ice concentration daily descending average		Int16	N/A	N/A
	Value	Description			
	0	Open Water			
	1 to 100	Percent Ice Concentration			
	120	Land Mask			
SI_25km_SH_ICEDIFF_ASC	Sea ice concentration daily ascending average using the difference between BBA and NT2 algorithms		Int16	N/A	N/A
	Value	Description			
	0	Open Water			
	1 to 100 or -1 to -100	Percent difference between algorithms*			
	120	Land Mask			
	200 to 300	Missing NT2 value (200+BBA)**			
	-200 to -300	Missing BBA value (-200-NT2)**			
-310	Out-of-range value in V12 data***				
SI_25km_SH_ICEDIFF_DAY	Sea ice concentration daily average using the difference between BBA and NT2 algorithms		Int16	N/A	N/A
	Value	Description			
	0	Open Water			
	1 to 100 or -1 to -100	Percent difference between algorithms*			
	120	Land Mask			
	200 to 300	Missing NT2 value (200+BBA)**			
	-200 to -300	Missing BBA value (-200-NT2)**			
-310	Out-of-range value in V12 data***				
SI_25km_SH_ICEDIFF_DSC	Sea ice concentration daily descending average using the difference between BBA and NT2 algorithms		Int16	N/A	N/A
	Value	Description			
	0	Open Water			
	1 to 100 or -1 to -100	Percent difference between algorithms*			
	120	Land Mask			
	200 to 300	Missing NT2 value (200+BBA)**			
	-200 to -300	Missing BBA value (-200-NT2)**			
-310	Out-of-range value in V12 data***				

Quality Assessment

A Quality Assessment (QA) XML metadata summary file is provided for each data file. The QA summary file denotes whether or not the file passed the science quality flag.

References

Cavalieri, D. and J. Comiso. 2000. Algorithm Theoretical Basis Document for the AMSR-E Sea Ice Algorithm, Revised December 1. Landover, Maryland USA: Goddard Space Flight Center.

Cavalieri, D. J., K. M. St. Germain, and C.T. Swift. 1995. Reduction of Weather Effects in the Calculation of Sea Ice Concentration with the DMSP SSM/I. *Journal of Glaciology* 41(139): 455-464.

Cavalieri, D. J., P. Gloersen, and W. J. Campbell. 1984. Determination of Sea Ice Parameters with the NIMBUS-7 SMMR. *Journal of Geophysical Research* 89(D4): 5355-5369.

Colbeck, S. C. 1982. An Overview of Seasonal Snow Metamorphism. *Reviews of Geophysics Space Physics* 20:45-61.

Comiso, J., D. Cavalieri, and T. Markus. 2003. Sea Ice Concentration, Ice Temperature, and Snow Depth Using AMSR-E Data. *IEEE Transactions on Geoscience and Remote Sensing* 41(2): 243-252.

Comiso, J. and K. Steffen. 2001. Studies of Antarctic Sea Ice Concentrations from Satellite Data and Their Applications. *Journal of Geophysical Research* 106(C12): 31,361-31,385.

Comiso, J. C., D. J. Cavalieri, C. L. Parkinson, and P. Gloersen. 1997. Passive Microwave Algorithms for Sea Ice Concentration - A Comparison of Two Techniques. *Remote Sensing of the Environment* 60: 357-384.

Comiso, J. C. 1995. SSM/I Ice Concentrations Using the Bootstrap Algorithm. NASA RP 1380.

Conway, D. 2002. Advanced Microwave Scanning Radiometer - EOS Quality Assurance Plan. Huntsville, AL: Global Hydrology and Climate Center.

Eppler, D. T. and 14 others. 1992. Passive Microwave Signatures of Sea Ice. IN: *Microwave Remote Sensing of Ice. Geophysical Monograph Series* 68: 47-71. Washington, D. C.: American Geophysical Union.

Fraser R. S., N. E. Gaut, E. C. Reifenstein, and H. Sievering. 1975. Interaction Mechanisms Within the Atmosphere Including the Manual of Remote Sensing. American Society of Photogrammetry 181-233. Falls Church, VA.

Gloersen P. and D. J. Cavalieri. 1986. Reduction of Weather Effects in the Calculation of Sea Ice Concentration from Microwave Radiances. *Journal of Geophysical Research* 91(C3): 3913-3919.

Kelly, R. E., A. T. C. Chang, L. Tsang, and J. L. Foster. 2003. A Prototype AMSR-E Global Snow Area and Snow Depth Algorithm. *IEEE Transactions Geoscience Remote Sensing* 41: 230-242.

Kummerow, C. 1993. On the Accuracy of the Eddington Approximation for Radiative Transfer in the Microwave Frequencies. *Journal of Geophysical Research* 98: 2757-2765.

Markus, Thorsten and Donald J. Cavalieri. 2008. [Supplement] AMSR-E Algorithm Theoretical Basis Document: Sea Ice Products. Greenbelt, Maryland USA: Goddard Space Flight Center. (PDF file, 2.10 MB)

Markus, T. and D. Cavalieri. 1998. Snow Depth Distribution over Sea Ice in the Southern Ocean from Satellite Passive Microwave Data. IN: *Antarctic Sea Ice: Physical Processes, Interactions, and Variability*. Antarctic Research Series 74: 19-39. Washington, DC: American Geophysical Union.

Markus, T., D. Cavalieri, and A. Ivanoff. 2011. Algorithm Theoretical Basis Document for the AMSR-E Sea Ice Algorithm, Revised December 2011. Landover, MD: Goddard Space Flight Center. (PDF file, 528 KB)

Markus, T. and D. Cavalieri. 2000. An Enhancement of the NASA Team Sea Ice Algorithm. *IEEE Transactions on Geoscience and Remote Sensing* 38: 1387-1398.

Matzler, C., R. O. Ramseier, and E. Svendsen. 1984. Polarization Effects in Sea-ice Signatures. *IEEE Journal of Oceanic Engineering* 9: 333-338.

Pearson, F. 1990. *Map projections: Theory and Applications*. Boca Raton, FL: CRC Press.

Snyder, J. P. 1987. *Map Projections - A Working Manual*. U.S. Geological Survey Professional Paper 1395. U.S. Government Printing Office. Washington, D.C.

Snyder, J. P. 1982. *Map Projections Used by the U.S. Geological Survey*. U.S. Geological Survey Bulletin 1532.

Worby, A. P., T. Markus, A. D. Steer, V. I. Lytle, and R. A. Massom. 2008. Evaluation of AMSR-E Snow Depth Product over East Antarctic Sea Ice Using In Situ Measurements and Aerial Photography, *Journal of Geophysical Research* 113(C05): Art. #C05S94. doi:10.1029/2007JC004181.

Contact Information

To order these data or for further information, please contact:

Global Hydrology Resource Center

User Services

320 Sparkman Drive

Huntsville, AL 35805

Phone: 256-961-7932

E-mail: support-ghrc@earthdata.nasa.gov

Web: <https://ghrc.nsstc.nasa.gov/>